



1. Start treatment at night before going to bed. Lying down will reduce leakage of the product from the vagina.
2. Wash the entire vaginal area with mild soap and water, and dry completely before applying the product.
3. *Vaginal cream*: (If prefilled applicators are being used, skip to step 4.) Unscrew the cap; place the cap upside down on the end of the tube. Push down firmly until the seal is broken. Attach the applicator to the tube by turning the applicator clockwise. Squeeze the tube from the bottom to force the cream into the applicator. Squeeze until the inside piece of the applicator is pushed out as far as possible and the applicator is completely filled with cream. Remove the applicator from the tube.
Vaginal tablets/suppositories: Remove the wrapper and place the product into the end of the applicator barrel.
4. While standing with your feet slightly apart and your knees bent, as shown in drawing A, or while lying on your back with your knees bent, as shown in drawing B, gently insert the applicator into the vagina as far as it will go comfortably.



A



B

5. Push the inside piece of the applicator in and place the cream as far back in the vagina as possible. To deposit vaginal tablets/suppositories, insert the applicator into the vagina and press the plunger until it stops.
6. Remove the applicator from the vagina.
7. After use, recap the tube (if using cream). Then clean the applicator by pulling the two pieces apart and washing them with soap and warm water.
8. If desired, wear a sanitary pad to absorb leakage of the vaginal antifungal. Do not use a tampon to absorb leakage.
9. Continue using the product for the length of time specified in the product instructions. Use the product every day without skipping any days, even during menstrual flow.